

$\Delta Q = 2$ Hyperon Decays

Ling-Fong Li

Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213

February 1, 2008

Abstract

The hyperon decays with 2 negative charged leptons in the final states can be of interest as possible tests for the Majorana neutrinos. The study of these decays are complementary to the double β decays of heavy nuclei.

It is now established that neutrinos do have non-zero masses[5][6][7]. The next step is to study the properties of these massive neutrinos. One interesting question is whether they are Majorana or Dirac neutrinos. Over the years the search for neutinoless double β decays of some heavy nuclei[1] is to test the possibility that neutrinos are Majorana particles. However, theoretical analysis for these processes is hampered by the complicate nuclear matrix elements which are very difficult to calculate reliably. As an alternative, we can study leptonic decays of the hyperons, $B_1 \rightarrow B_2 ll'$. Here l and l' are same sign leptons, like e^-e^- , $\mu^-\mu^-$, or $e^-\mu^-$. The study of these processes can serve as complement to the nuclear double β decays. Some upper bounds already exist for some of these processes[2][3].

We can classify the possible hyperon decays by the change in the strangeness.

1. $\Delta S = 0$

$$\Sigma^- \rightarrow \Sigma^+ e^- e^-, \quad M(\Sigma^-) - M(\Sigma^+) = 8MeV$$

2. $\Delta S = 1$

$$\Sigma^- \rightarrow p e^- e^-, \quad M(\Sigma^-) - M(p) = 257 \text{ MeV}$$

$$\Sigma^- \rightarrow p \mu^- e^-, \quad M(\Sigma^-) - M(p) - M(\mu) = 151 \text{ MeV}$$

$$\Sigma^- \rightarrow p \mu^- \mu^-, \quad M(\Sigma^-) - M(p) - 2M(\mu) = 46 \text{ MeV}$$

$$\Xi^- \rightarrow \Sigma^+ e^- e^-, \quad M(\Xi^-) - M(\Sigma^+) = 132 \text{ MeV}$$

$$\Xi^- \rightarrow \Sigma^+ \mu^- e^-, \quad M(\Xi^-) - M(\Sigma^+) - M(\mu) = 26 \text{ MeV}$$

$\Delta S = 2$

$$\Xi^- \rightarrow p e^- e^-, \quad M(\Xi^-) - M(p) = 383 \text{ MeV}$$

$$\Xi^- \rightarrow p \mu^- e^-, \quad M(\Xi^-) - M(p) - M(\mu) = 277 \text{ MeV}$$

$$\Xi^- \rightarrow p \mu^- \mu^-, \quad M(\Xi^-) - M(p) - 2M(\mu) = 172 \text{ MeV}$$

$$\Omega^- \rightarrow \Sigma^+ e^- e^-, \quad M(\Omega^-) - M(\Sigma^+) = 483 \text{ MeV}$$

$$\Omega^- \rightarrow \Sigma^+ \mu^- e^-, \quad M(\Omega^-) - M(\Sigma^+) - M(\mu) = 377 \text{ MeV}$$

$$\Omega^- \rightarrow \Sigma^+ \mu^- \mu^-, \quad M(\Omega^-) - M(\Sigma^+) - 2M(\mu) = 272 \text{ MeV}$$

We see that the $\Delta S = 0$ decay $\Sigma^- \rightarrow \Sigma^+ e^- e^-$ has very small phase space to be of interest and there are many other processes which have comparable phase space as the usual hyperon leptonic decays.

Effective interactions

The Majorana neutrinos can give rise to the decay processes discussed here just like the case in the nuclear double β decays, e.g.

But there can also be some new interaction where a new doubly charged scalar particle couples to dileptons as shown in the figure[4],

Of course there are other mechanism for these decays and they will undoubtedly involve new particles and/or new interactions. Thus the study of these processes can shed some light on either Majorana nature of the neutrinos or uncover some new physics beyond the standard model. Here we will just discuss some general features of these processes without going into details of each possible models. It is reasonable to assume the effective interaction for

the processes of interest can be written as

$$L_{\beta\beta} = \frac{G_F^2}{\Lambda_{\beta\beta}} [c_1(\bar{u}\Gamma_i d)(\bar{u}\Gamma_j d) + c_2(\bar{u}\Gamma_i d)(\bar{u}\Gamma_j s) + c_3(\bar{u}\Gamma_i s)(\bar{u}\Gamma_j s)] \\ \times [d_1(\bar{e}\Gamma_k e^c) + d_2(\bar{e}\Gamma_k e^c) + d_3(\bar{e}\Gamma_k \mu^c + \bar{\mu}\Gamma_k e^c)]$$

Here $\Lambda_{\beta\beta}$ is a parameter which characterizes the physics scale responsible for these processes and c_i, d_i are dimensionless parameters which represent different the interaction strengths for various channels. These parameters depend on specific model and certainly will contain some unknown parameters. Also Γ_i 's are some combinations of Dirac gamma matrices which depend on the physical mechanism involved.

Since the matrix elements of the 4-quark operators are difficult to calculate, we will just get some order of magnitude estimate of these decays. The hadronic matrix elements of 4-quark operator will be compared to that of the hyperon leptonic decays,

$$R_h = \frac{\langle B_2 | (\bar{u}\Gamma_i d)(\bar{u}\Gamma_j s) | B_1 \rangle}{\langle B_2 | ((\bar{u}\Gamma s) | B_1 \rangle}$$

This ratio has mass dimension 3 and we will write it as

$$R_h = M^3$$

where M is some mass parameter. It is possible that M is of order of hadron mass of 1 *Gev* or so. We can write the decay rates as

$$\frac{\Gamma(B_1 \rightarrow B_2 l l')}{\Gamma(B_1 \rightarrow B_2 e \nu)} \sim \left(\frac{G_F}{\Lambda_{\beta\beta}} \right)^2 M^6 \rho$$

where ρ represents the phase spaces ratio. If we take M to be of order of 1

Gev or so, this ratio is of order of

$$\frac{\Gamma(B_1 \rightarrow B_2 l l')}{\Gamma(B_1 \rightarrow B_2 e \nu)} \sim \left(\frac{M}{\Lambda_{\beta\beta}} \right)^2 \rho \times 10^{-10}$$

In the usual hyperon leptonic decays, when the lepton masses are neglected, the phase space is proportional to

$$\Delta^5 = (M_1 - M_2)^5$$

where M_1, M_2 are masses of B_1, B_2 respectively. This should apply also to decay channels with 2 electrons in the final states. But for cases where there are muons in the final states Δ should be replaced by

$$q = \sqrt{(M_1 - M_2)^2 - m^2} \quad \text{or} \quad \sqrt{(M_1 - M_2)^2 - 4m^2}$$

where m is the muon mass. In most cases, the phase space does not give large suppression or enhancement except for the decay $\Sigma^- \rightarrow \Sigma^+ e^- e^-$ where the phase space suppression is of order of 10^{-8} or so. In some cases with $\Delta S = 2$, there is a small enhancement due to the fact the mass differences can be larger than those with $\Delta S = 1$. For example, in the decay $\Xi^- \rightarrow p e^- e^-$ the phase space is larger by a factor of 200 over that of $\Xi^- \rightarrow \Sigma^0 e^- \nu_e$.

For the case of Majorana neutrinos, the factor $\frac{1}{\Lambda_{\beta\beta}}$ should contain a factor of Majorana mass of neutrinos m_{maj} ,

$$\frac{1}{\Lambda_{\beta\beta}} = \left(\frac{m_{maj}}{\Lambda'} \right) \frac{1}{\Lambda'}$$

and will probably give a large suppression factor. Thus branching ratio of order of 10^{-10} is probably most optimistic upper bound for these type of decays and most likely they are much smaller than this.

References

- [1] For a review, see for example, Petr Vogel, "Neutrinoless double beta decay", To appear in the proceedings of Exploring New Frontiers Using Colliders and Neutrinos (TASI 2006), Boulder, Colorado, 4-30 Jun 2006. e-Print: hep-ph/0611243
- [2] Laurence S. Littenberg and Robert Shrock, "Upper bounds on $|\Delta L| = 2$ decays of baryons", Phys. Rev. **D46**, 892 (1992)
- [3] D. Rajaram et al " Search for the Lepton-Number-Violating Decay $\Xi^- \rightarrow p\mu^-\mu^-$ ", Phys Rev Lett **94**, 181801(2005).
- [4] See for example, T.P. Cheng and Ling-Fong Li, " Neutrino masses, mixings, and oscillations in $SU(2) \times U(1)$ models of electroweak interactions", Phys. Rev. **D22**, 2860,(1980)
- [5] SuperKamiokande Collaboration, Y. Fukuda et al., Phys. Rev. Lett. 81, 1158 (1998); 81, 1562 (1998); 82, 2644 (1999); 85, 3999 (2000); Y. Suzuki, Nucl. Phys. B, Proc. Suppl. 77, 35 (1999); S. Fukuda et al., Phys. Rev. Lett. 86, 5651 (2001); Y. Ashie et al., Phys. Rev. Lett. 93, 101801 (2004).
- [6] KamLAND Collaboration, K. Eguchi et al., Phys. Rev. Lett. 90, 021802 (2003); T. Araki et al., Phys. Rev. Lett. 94, 081801(2005).
- [7] SNO Collaboration, Q. R. Ahmad et al., Phys. Rev. Lett. 89, 011301 (2002); 89, 011302 (2002); 92, 181301 (2004); B.Aharmim et al., Phys. Rev. C 72, 055502 (2005).